

## Faunistic and taxonomic notes upon some histerids (Coleoptera, Histeridae)

Slawomir Mazur

Mazur S. 2010. Faunistic and taxonomic notes upon some histerids (Coleoptera, Histeridae). *Baltic J. Coleopterol.*, 10(2): 141- 146.

New records for particular species are given. *Platysoma celatum* has been transferred to the genus Kanaarister. The male genital structure is first time figured for *Hister capicola*, *H. castus* and *Eugrammicus proximus*, for *Campylorhabdus poggei* some morphological details are figured. The taxonomic status of *Hister castus* is discussed. *Eupoachylopus ripae* ab. *therskii* is synonymized with the nominative form. *Idister modigliani* is a distinct species.

Key words: faunistics, taxonomy, Histeridae, new records

Slawomir Mazur; Department of Forest Protection and Ecology, WULS, Nowoursynowska 159, bld. 34, 02-776 Warszawa, Poland; e-mail: slawomir.mazur@wl.sggw.pl

### INTRODUCTION

This paper is a result of the elaboration of the own author's collection (CHSM). Some materials of the collection of Tomas Lackner were also used. Additionally, the results of the studies upon the type materials kept at the Museum of Natural History, London (UK) [MNHL] and the Muséum National d'Histoire naturelle, Paris (France) [MNHN] were presented here.

### ANALYTICAL PART

*Baconia venusta* (LeConte, 1845)

Mexico: Chetumal, Quintana Roo, 7.10.2004, 3 exs., leg. M. Sawoniewicz [CHSM].

Known from the southern states of the USA. New to Mexico.

*Nunbergia exosternoides* Mazur, 1978

Argentina: Neo – Chaco – Charata Dept., Chacabuco – Pampa Avila, Nest von Schleiereule

- *Tyto alba* (Scop.), 21.09.1995, 1male, 1female, leg. Liebig [CHSM]. Known only from Brazil. New to Argentina.

*Apobletes platessae* Lewis, 1900

RSA: Natal, 1 ex., leg. Dr. Martin [CHSM]. Rarely collected, known from Tanzania. New to RSA.

*Platysoma inexpectatum* Lackner, 2004

Turkey: Mersin, 40 km N Gülnar, N 36°30'22.5"; E 33°07'43.3", Kösecabanli/Tasdüstü, hollow *Quercus*, 24-06-2006, 2 exs, leg. N. Jansson & M. Avci [CHSM].

Described and recorded only from Syria as yet. New to Turkey.

*Platysoma (Cylister) yunnanum* (Kryzhanovskij, 1972)

China: Zhejiang prov., Lin'an County, W. Tianmu Shan N.R., 1000 m, 18.V.1996, 1 ex., leg. J. Cooter [CHSM].

Described and recorded only from Yunnan. New locality.

*Kanaarister celatus* (Lewis, 1884) comb. nov.  
China: Guizhou prov., 20 km NW of Jiangkou, Fanjing Shan – Kuaichang, 27.V-3.VI.1995, 1 ex., leg. E. Jendek & O. Sausa [CHSM].

Recorded from Japan, Taiwan and south-eastern China: Guandong, Zhejiang.

Note. Because of the sternal structure (short, widely sinuous mesosternum, absence of the post mesocoxal striae) this species should be transferred from *Platysoma* Leach to *Kanaarister* Mazur.

*Idister modigliani* Schmidt, 1894, bona sp.  
Malaysia: Pahang distr., 30 km NE Raub, Lata Lembik, 3°56' N, 101°38' E, 200–400 m, 22.IV.–1.V.2002, 3 exs, leg. E. Jendek & O. Sausa [CHSM, CHTL].

Described over a hundred years ago but still known only from the type locality: Indonesia, Palau Enggano Is. New to Malaysia.

Note. Considering almost same distribution of all the species Mazur (1997: 26) synonymized *I. modigliani* and *I. mendax* Lewis with *I. morphon* Marseul. In the next studies dealing with the genus *Idister* Marseul (Mazur & Ôhara, 2006: 35–36) the identity of *I. mendax* has been fully confirmed. Finding of the next specimens of *I. modigliani* in Malaysia allowed to state it is a distinct species.

*Hololepta nepalensis* Lewis, 1910  
China: Fujian prov., Wuyi Shan Nat. Res., Quiligao – Guandan Road, 1000–1330 m, 1.06.2001, under bark of dead tree, 1 ex., leg. Hlavac & Cooter [CHSM].

Poorly known species, recorded from Nepal, Vietnam and Indonesia: Borneo. New to China.

*Scapomegas gibbus* Marseul, 1855  
Paraguay: Colonia Indigena Chupa Pou, 11 km S of Ygatimi Dept., Canindeyu, 20-23.XI.1994, carcass bait trap, 1 ex., leg. B. Malkin; Dep. Alto Parana, Estancia Dimas, S 25°32.6', W 55°13.4', 17-22.02.2005, 1 ex., leg. U. Dreschel [CHSM].  
A rare species, known only from Brazil as yet. New to Paraguay.

#### *Hister capicola* Marseul, 1854

A rare species, endemic to South Africa. To a better recognition the figures of the male genital structure of the type are given here (Figs. 1-5). Type material. Male, 37, *Hister capicola*, Cap.[land], Museum Paris, col. de Marseul, 2842–90 [MNHN].

Other material studied. Südafrika, Somerset, West Cape Proviru, 28.10.1990, 1 ex., leg. A. Taubert [CHSM].

#### *Hister castus* Lewis, 1885

This species belongs to a specific species-group, including *H. castus*, *H. honestus* Lewis, 1908 and *H. martius* Lewis, 1885. After having studied the type of *H. honestus* the author (Mazur, 2007: 139) followed the Bickhardt's supposition that this species might have been belong to the genus *Eudiplister* Reitter, 1909. Bickhardt (1919: 84) also suggested that all these three species were identical. To resolve this problem the author also studied the types of *H. castus* and *H. martius*, finding any substantial external differences between them. Unfortunately, only the type of *H. castus* was a male (one male was also studied by Penati & Vienna, 2006: 24), so now it is almost impossible to confirm the identity of the species mentioned.

Having most of the external characters of *Eudiplister*; *H. castus* differs, however, from the Palearctic representatives of the genus *Eudiplister* by a different structure of the maleedeagus, especially in ratio of basal piece to parameres (Figs. 6-11).

If this character is confirmed on larger materials, there will be, probably, a need to create a new genus for these three (or only one?) species.


Type material.

*Hister castus*: Male, [Saudi Arabia] Arabia, El Hejaz [El Hahaz], [leg.] Millingen, *Hister castus* Lewis, Type, G. Lewis Coll., B.M. 1926-369 [MNHL].

*Hister martius*: Female, [Ethiopia] Abyssinia, [leg.] Raffray, *Hister martius* Lewis, Type, G. Lewis Coll., B.M. 1926-369 [MNHL].

Other material studied. [Malawi] Nyassa-See, Kigonsera, 1 ex. [CHSM].

#### *Campylorrhynchus poggei* (Harold, 1878)



Figs. 1-11. Genital structure of the male. 1-5 – *Hister capicola*, 6-9 – *H. castus*, 10, 11 – *Eudiplaster planulus*. 1-3, 6-8, 10, 11 – edeagus, 4, 9 – 8<sup>th</sup> segment, 5 – 9<sup>th</sup> and 10<sup>th</sup> tergites, 1, 6 – dorsally, 3, 7, 11 – ventrally, 2, 8, 10 – laterally.

The genital male genital structure of this rare species was presented earlier (Mazur, 2008: 45, figs. 21-25). Additionally, some details of the external morphology are figured here (Figs. 12-15).


Material studied. Malawi, Chitipa Distr., Jembya reserve, 18 km SSE Chisenga, 10-08 S, 33-27 E, 1870 m, 1-10 Jan. 1989, 1 ex., leg. J. Rawlins & S. Thompson; Zambia, 8-9.12.2004, 1 ex., leg. K. Werner [CHSM].

#### *Eugrammicus proximus* (Lewis, 1906)

A rare African species, one of two representatives of the genus, the genital structure for which has never been studied before (Figs. 16-19).

Type material. Male, Niger, Warri, March [18]96, Dr. Roth, Contipus proximus Lewis, Type, George Lewis Coll., B.M. 1926-369 [MNHL].

Other material studied. [Tanzania], Uamgebiet, Bosum, 21-30.4.[19]14, 1 ex., leg. S. Tessmann [CHSM].



Figs. 12-19. 12-15 – *Campylorrhynchus poggei*. 16-19 – *Eugrammus proximus*. 12 – upper side, 13 – body, laterally, 14 – pro- and pygidium, 15 – under side, 16, 17 – edeagus, 18 – 8<sup>th</sup> segment, 19 – 9<sup>th</sup> and 10<sup>th</sup> tergites, 16 – ventrally, 17 – laterally

<i>Nagelius carinicollis</i> (Lewis, 1893) Laos: Bolikhamsai prov., Ban Nape, 16.07.2006, 18°20'N, 105°E, 1 ex., leg. Jendek & Sausa [CHSM]. Rare species, known only from Borneo. New to Laos.	China: Tibet, South-west Himalaya Mts., Dongpa, 4500 m, 11.VII.1997, 3 exs, leg. L. Bieber [CHSM]. A poorly known species, described and recorded so far only from Nepal. New to China.
<i>Diplostix vicaria</i> (Cooman, 1935) Taiwan: Nantou Hsien, NTU Exp. Forest, <i>Cunninghamia lanceolata</i> , Nei-mou-pu Trap, pheromon trap, 26.07.2004, 1 ex., [CHSM]. Recorded from Indochina (Vietnam, Thailand, Laos), Indonesia (Java) and China (Yunnan). New to Taiwan.	<i>Eopachylopus ripae</i> (Lewis, 1885) China mer., Hongkong, 1993, 1 ex., leg. G. de Rougemont [CHSM]. Known from Russia : Primorskiy Kray, South Korea and Japan. New to China. Note. When describing Lewis (1885: 469) noted that it occurred in two color forms: "Some example are unicolorous, others have a red band across the middle of the elytra". The last form was described first by Reichardt (1941: 407) as ab. <i>therskii</i> Reichardt and later by Osawa (1952: 4) as f. <i>rufofasciatus</i> Osawa. The last name has been synonymized with <i>Eopachylopus ripae</i> by Mazur (2004: 24). Thus, <i>Eopachylopus ripae</i> ab. <i>therskii</i> Reichardt, syn. nov. of <i>E. ripae</i> (Lewis).
<i>Chaetabraeus controversus</i> (Cooman, 1935) Laos: Bolikhamsai prov., Bon Nape, 16.07.2006, 18°20' N, 105°E, 3 exs, leg. E. Jendek & O. Sausa [CHSM, CHTL]. Rarely collected, known from Vietnam and Thailand. New to Laos.	
<i>Plegaderus vulneratus</i> (Panzer, 1797) China: Heilongjiang prov., Quing Yuan, ca 30 km S Lang Xian, 46°47.470' N, 129°03.823' E, ca 600-700 m, 25-29.V.2005, 2 exs, leg. J. Cooter [CHSM]. Forest zone of the Palearctic. New to China.	
<i>Acritus nepalensis</i> Gomy, 1976 India: Uttar Pradesh, 2 km E de Dhamolti, 2250 m, 21.X.1979, 1 ex., Kumaon, Rangarh, env. 2000 m, 9.X.1979, 1 ex., leg. I. Löbl [CHSM]. Nepal. New to India.	<b>ACKNOWLEDGEMENTS</b> The author is deeply indebted to Dr. Roger Booth (MNHL) and Dr Azadeh Taghavian (MNHN) for the privilege of studying the types of G. Lewis and S.A. Marseul. Special thanks are due to Tomas Lackner (The Hokkaido University Museum, Sapporo, Japan) for having enabled the studies upon his histerids.
<i>Acritus punctisternus</i> Wenzel & Dybas, 1941 Panama: Barro Colorado, 5 exs., Gamboa, 5 exs., leg. K. Burchard [CHSM]. Described and recorded only from Colombia. New to Panama.	<b>REFERENCES</b> Bickhardt H. 1919. Die Histerini des aethiopischen Faunengebiets (Col, Histeridae) (41. Beitrag zur Kenntnis der Histeriden). Abhandlungen und Bericht des Vereins für Naturkunde zu Cassel. 81-83(1916-1919): 1-158. Lewis G. 1885. New species of Histeridae, with synonymous notes. The Annals of Natural History. 5(15): 456-473. Mazur S. 1997. A world catalogue of the Histeridae. Genus, Suppl. 1997, 373 pp. Mazur S. 2004. Histeridae, pp. 68-102. In: I. Löbl & A. Smetana (editors): Catalogue of
<i>Acritus tuberisternus</i> Cooman, 1932 Japan: Ryukyus, Kuroshima, 8.12.2008, under bark, 2 exs, leg. S. Mazur & M. Ôhara [CHSM]. Widely distributed in the Indo-Pacific region (Mauritius, Sri Lanka, India, Myanmar, Vietnam, Laos, Malaysia, Taiwan). New to Japan.	
<i>Saprinus subcoeruleus</i> Thérond, 1978	

---

Received: 18.11.2010.

Accepted: 15.12.2010.

- Palaearctic Coleoptera, Vol. 2, Stenstrup:  
Apolo Books, 942 pp
- Mazur S. 2007. Notes on some African Histerini  
(Coleoptera: Histeridae) with description of  
a new genus Quassarus gen. n. Studies and  
reports of District Museum Prague-East,  
Taxonomical Series. 3(1-2): 129-140.
- Mazur S. 2008. Notes on some African Histerini  
(Coleoptera: Histeridae). Baltic Journal of  
Coleopterology. 8(1): 41-47.
- Mazur S., Ôhara M. 2006. Notes on the genera  
related to Idister Marseul, 1880 (Coleoptera,  
Histeridae). Annals of Warsaw Agricultural  
University – SGGW, Forestry and Wood  
Technology, 60: 35-38.
- Osawa S. 1952. On some species of Histeridae  
from Japan and its adjacent district (Col.).  
The Entomological Review of Japan. 6: 4-6.
- Penati F., Vienna P. 2006. An updated catalogue  
of the Histeridae (Insecta: Coleoptera) of the  
Arabian Peninsula, with biogeographical  
remarks. Zootaxa, 1157: 1-74.
- Reichardt A. 1941. Sem. Sphaeritidae I Histeridae  
(chast' 1). In: Fauna SSSR, Nesekomye  
zhestkokrylye, V. 3. Moskva-Leningrad: Izd.  
Akademii Nauk SSSR, xiii+419 pp.